Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.





Ag 8 United States Department of Agriculture

Nutrition Information Service

Home Economics Research Report Number 45

Iron Content of Food

Jacob Exler

IRON CONTENT OF FOOD, Jacob Exler Consumer Nutrition Center, Human Nutrition Information Service, U.S. Department of Agriculture, Hyattsville, Maryland 20782. Home Economics Research Report Number 45

ABSTRACT

This publication presents newly compiled compositional data on the iron content of 277 foods commonly eaten in this country. Values are given only for food groups for which sections of revised Agriculture Handbook No. 8 have not yet been published. This provisional table will be superseded by subsequent sections of revised Handbook No. 8 as they are published.

The level of reliability of the values is given by the number of observations on which the mean values were based and the corresponding sample standard errors. Another measure of reliability, included in a table of food composition for the first time, is the confidence code. This code, based on a critical evaluation of the sources of data from which the mean values were derived, gives an indication of the confidence one can have in how closely the iron content of a food item is represented by the published value.

KEYWORDS: Iron content, foods, reliability, quality index, confidence code.

Washington, D.C.

ACKNOWLEDGMENTS The author gratefully acknowledges Susan Akridge for her technical assistance; the food group specialists of the Nutrient Data Research Group, Consumer Nutrition Center, for providing their collected data; and Wayne Wolf and Mary Moss, Nutrient Composition Laboratory, Beltsville Human Nutrition Research Center, Kent Stewart, Virginia Polytechnic Institute and State University, and Frank Hepburn, Nutrient Data Research Group, for their critical discussions in the development of the confidence codes and alternate means of nutrient data evaluation.

		Page
CONTENTS	INTRODUCTION	1
	RELIABILITY OF VALUES	2
	DESCRIPTION OF THE TABLE	3
	IRON BIOAVAILABILITY	4
	Tables 1 Criteria for quality index	2 3 5 5 6 6 9 10 10 11 12 12 12
	I.TTERATURE CTTED	15

Iron Content of Food

Jacob Exler¹

INTRODUCTION

Many people in the United States have dietary intakes of iron, as estimated by using current food consumption tables, which are below the Recommended Dietary Allowance (RDA) $(\underline{1},\underline{2})^2$. To improve estimates of intakes and provide the basis for dietary guidance, more reliable data on the iron content of food are necessary.

This provisional table updates many of the iron values in Agriculture Handbook No. 8 (3) and in other U.S. Department of Agriculture publications by presenting newly compiled compositional data on the total iron content of foods commonly eaten in this country. Some of the earlier data were based on older, less reliable methods and some were based on relatively few samples. Also, because of the lack of sufficient analytical data, previously published values for iron in most meat products had been calculated as a function of protein as discussed in Appendix B of Agriculture Handbook No. 8 (3).

Iron values are given here for items in food groups for which sections of revised Agriculture Handbook No. 8 have not yet been published. These values were obtained from summaries compiled by the staff at the Consumer Nutrition Center, from data on file and from the USDA Nutrient Data Bank. The values reported here are summarized from data obtained by using improved, iron-specific analytical methods. Sources of data were the published literature, government and academic publications, and industrial and analytical laboratory reports.

For the iron content of items in those food groups for which sections of Handbook No. 8 have been revised and published [Dairy and Egg Products ($\underline{4}$), Spices and Herbs ($\underline{5}$), Baby Foods ($\underline{6}$), Fats and Oils ($\underline{7}$), Poultry Products ($\underline{8}$), Soups, Sauces, and Gravies ($\underline{9}$), and Sausages and Luncheon Meats ($\underline{10}$), and Breakfast Cereals ($\underline{11}$)], the appropriate sections should be consulted 3 .

This provisional table will be superseded by subsequent sections of revised Handbook No. 8 as they are published. The data in these future handbook sections will incorporate additional data received after preparation of this table. Therefore, some revision and additions to the iron values presented here should be anticipated in the future sections.

¹The author is a Research Chemist with the Consumer Nutrition Center, Human Nutrition Information Service.

²Underlined numbers in parentheses refer to Literature Cited, p. 5.

³The revised section on Fruits and Fruit Juices (AH-8-9) was published while this report was in press.

RELIABILITY OF VALUES

Many factors affect the iron content of foods. Content may vary with the time and location of harvest of the raw food. Some processing procedures remove iron from the food while others inadvertently add the nutrient by contamination. Differences in product formulation among manufacturers are another source of iron variability.

The available data and information for iron in the foods listed were not equally satisfactory for deriving values. The level of reliability of the values published in this table is indicated in several ways. Shown are the number of observations on which the mean value is based and the corresponding sample standard error (if the number of samples is greater than two). Also included is a confidence code.

Confidence codes are for use in food tables and in provisional nutrient tables. The purpose of these codes is to give a user an indication of the confidence he can have in the mean value given in the table. The codes are based on a critical evaluation of the data sets from which the mean values were derived. A quality index (0, 1, 2, or 3) for the data from each data set is assigned by using the criteria in table 1.

TABLE 1.--CRITERIA FOR QUALITY INDEX

Evaluation	Documentation of analytical method	Sample handling and appropriateness of analytical method	Quality control
0	None	Totally incorrect handling	No dupli- cates
1	Unpublished, but written	No documentation	Duplicate aliquots
2	Published, but modified	Reasonable, docu- mented common technique	Duplicate samples
3	Complete pub- lished writeup	Extensive docu- mented testing and appropriate method was used	Standard reference materials, spikes, recoveries, or blind duplicates

The lowest value for any criteria becomes the limiting quality index for the data from each data set. The sum of the quality indices from all the data sets used to obtain the overall mean value is the basis for selecting the confidence code for that mean value.

Sum of quality indices	Confidence code	Meaning of confidence code
<u>></u> 6	a	The user can have confience in the mean value.
3-5	Ъ	The user can have some confidence in the mean value; however, some questions have been raised about the value or the way it was obtained.
1-2	с	There have been some serious questions raised about this value. It should be considered only as a best estimate of the level of this nutrient in this food.

The values reported in the table are the means of the data from two or more sources of data in which the mean values from each source do not differ from the overall mean by more than 30 percent of the overall mean. Other data are designated by an asterisk. A single asterisk (*) denotes that the data are from a single source. Two asterisks (**) denote that the data are from two or more sources, but the means differ from the overall means by more than 30 percent of the overall mean.

The data presented in this table, and in food composition tables in general, are intended to represent values of the nutrient content of food on a nationwide, year-round basis. The information on the reliability of each value in this table should be used to assess the confidence in how closely the iron content of a food sample is represented by that value.

DESCRIPTION OF THE TABLE

In this table, each food item is described and the mean iron content in milligrams per 100 grams of the edible portion of the food is given. The sample standard error and the total number of observations are also given. Next are the confidence code and asterisk(s), where appropriate. In the last column, the item numbers are given for those foods that correspond to items in the 1963 edition of Handbook No. 8. Dashes in this column denote there was no corresponding food or no value for iron was given in Handbook No. 8.

In general, foods that provided less than 2 percent of the U.S. Recommended Daily Allowance (U.S. RDA) for iron per serving were not included in the table. This corresponds to 0.36 milligram per serving and is the level below which, according to Federal

regulations $(\underline{12})$, no specific value need be included on a nutrient label. Some foods with iron content below 0.36 milligram per serving were included in the table because values differ from these previously reported, because the food is consumed frequently or in large quantities, or to provide comparison with a closely related food.

"Enriched," as used in this table, refers only to iron and not to any other added nutrient. Foods with added iron must have the iron content per serving listed on the label expressed as the percent of the U.S. RDA. The following chart relating the percent of the U.S. RDA to milligrams is included to assist in estimating iron content from label claims:

Perc	ent																										
U.S.	RDA																										Iron
																											(<u>mg</u>)
2																											0.36
2	• • •	• •	• •	-	 -	-	 -	-	-	-	-	-	-	-	-	-	-	-	_	 -	-	-	-	-	-	-	0.30
4		• •	• •	•		•	 •	•	•	• •	•	•	•	• •		•	•	•	•	•	•		•	•	•	•	.72
8					 •	•		•			•		•			•		•	0 (•	•	•		1.4
10																			•								1.8
15		• •						•																			2.7
25						•							•				•		•								4.5
45		• •				•							•			•			• (•	•			8.1
50																			•								9.0
100		• •	• •			•					٠						•	•	• (٠							18.0

Actual iron content may be higher than that claimed on the label as a consequence of the compliance regulation $(\underline{12})$.

IRON BIOAVAILABILITY

The values in this table are for the total iron content of the food items and do not take into consideration the bioavailability of the iron in the foods. An estimate of the available iron in a meal can be calculated by the method of Monsen et al. (2, 13).

Item No.	Food	Amount Mean	of iron in Standard error	100 grams Number of samples	Confi- dence code ¹	AH-8 Item No. (1963)
<u></u>		Milligrams				
	BAKERY PRODUCTS					
	Breads:					
1	Cracked wheat	2.6	0.42	4	b**	444
2	French, enriched	2.8	.12	38	a	446
3	Mixed grain	3.2	.09	136	a**	
4	Raisin	2.9	.29	11	Ъ	452
5	Pumpernickel	2.9	.19	4	Ъ	456
6	Regular	2.7	.10	43	Ъ	454
7	Wheat	3.5	•05	140	Ъ	
8	White, enriched	3.0	.02	445	a	461
9	Whole wheat	3.2	.15	27	а	471
10	Danish pastry	1.8	.10	9	Ъ	1899
11	English muffins, plain	2.8	.09	25	a**	
12 13	Dinner, enriched	3.1	.07	110	a	1902
	enriched	3.0	.03	250	a	1902
14	Rye	2.8	()	2	Ъ	
15	Tortillas, corn	1.9	.06	6	C*	
	BEEF					
16	Hamburger, lean, cooked	2.7	.16	4	b*	368
17	Lean meat, cooked	2.7	.08	79	b*	
18	Liver, fried	5.7	1.2	5	b*	1267
	BEVERAGES					
	Alcoholic:					
19	Beer	.01	.001	66	a**	394
20	Red	.94	.03	172	a**	401
21	White	•57		208	a**	401
22	Carbonated, nonalcoholic	.07		113	c*	402
	our point con in the contract of the contract	•••	V 0 =			to 409
	Chocolate flavored beverage powders (add milk):					407
23	Plain	3.4	.31	19	a**	779
24	With added nutrients	21.5	2.0	3	a	
25	Cocoa mix powder (add water)	1.1	.11	9	a	780
	TITUS MEN PONGOL (add Nacol) *****	1 • 1	***		u	, 00

Continued--

5

Item No.	Food	<u>Amount</u> Mean	of iron in Standard error	100 grams Number of samples	Confi- dence code ¹	AH-8 Item No. (1963)
		Milligrams				
	Coffee:					
26	Instant, powder	3.7	0.51	5	a**	799
27	Ground, brewed	.01	(²)	1	b*	
28	Tea, instant, powder: Sweetened	.15	•01	3	b*	
		2.0	(2)	1	L. *	2276
29	Unsweetened	3.8	(²)	1	b*	2270
	FISH PRODUCTS ³					
	Crustaceans:					
	Raw meat:	F.0	.07	12	Ъ	
30	Crab, blue	.58 1.8	.42	6	b*	2042
31	Shrimp	1.0		U	_	
32	Shrimp	1.7	.38	6	b*	2045
	Finfishes:					
2.2	Raw fillet:	.84	(²)	1	C*	
33 34	Bass, striped	.43		3	b**	794
35	Flatfishes	.31		13	a	1018
36	Ocean perch, Atlantic	.92		24	b**	1396
37	Salmon, sockeye	.64	2	2	b *	
57	Canned:		()	_		
	Solids:					
38	Herring, Atlantic, in oil	3.1	.34	4	c*	
39	Salmon, sockeye, in water	.78	.05	20	a**	
4.0	Solids and liquid:	.83	.04	25	b*	1955
40	Salmon, pink, in water Sardines, in tomato sauce	2.4	.09	29	Ъ	1976
41	Tuna, in water or oil:	2.4	•03	23	В	1370
42	Light meat	1.5	.05	107	a**	
43	White meat	.56	.06	35	a	
	Mollusks, raw meat:					
44	Clam, hard shell	3.0	.15	106	Ъ	771
45	Oyster, Eastern	6.5	.29	164	Ъ	1443
	FRUITS AND FRUIT JUICES3					
	Apples:					
46	Raw	.18		119	a**	13
47	Canned, sliced	.23		6	b*	
48	Dried	1.4	.10	37	a**	21
49	Juice, canned	•37	.03	11	b**	27
					Cont	inued

TABLE 3.--IRON CONTENT OF EDIBLE PORTION OF FOOD--Continued

Item No.	Food	Amount Mean	of iron in Standard error	100 grams Number of samples	Confi- dence code ¹	AH-8 Item No. (1963)
		Milligrams				
50	Applesauce, canned, sweetened Apricots:	0.35	0.03	43	a**	29
51	Raw	•54	.08	31	a**	30
52	Water pack	.54	(²)	1	b*	32
53	Juice pack	.30		8	a**	33
54	Light	.39	.03	37	b*	34
55	Heavy	.30		156	a	35
56	Dried, sulfured	4.7	.18	26	a**	39
57	Nectar, canned	•38	.02	68	a**	43
58	Avocados, raw	1.0	.12	54	b**	64
59	Bananas, raw	.31	.02	108	a**	141
60	Blackberries, raw	.44		16	c*	417
61	Raw	.16	.01	114	a**	424
62	Frozen, sweetened	.39		3	c*	428
63	Cherries, sour, red, raw	.32		347	a**	662
64	RawCanned:	.39	.02	57	a**	663
65	Water pack	.36	(²)	1	c*	669
66	Juice pack	.58	.12	6	C*	
67	Sirup pack, heavy	.35	.01	82	b*	671
68	Frozen, sweetened	.35	•04	3	c *	
69	Raw	.20	.02	16	c*	920
70	Canned, jellied sauce	.22	(²)	2	c*	923
71	Juice drink	.16	.01	4	c*	922
72	Raw	1.2	•23	8	а	944, 945
73	Dried, Zante	3.2	.16	37	а	
74	Dates, dried, pitted	1.2	.19	17	b**	952
75	Canned, sirup pack, heavy	.28	.02	18	b*	1005
76	DriedFruit cocktail, canned:	2.2	•16	36	a**	1007
77	Water pack	.25	.03	13	b**	1021
78	Juice pack	.21	.03	7	b**	
79	Sirup pack, heavy	.29	.01	160	а	1023
80	Juice pack	.25	.09	7	b**	
81	Sirup pack, heavy	.28	.01	92	a	1027

Continued--

Item No.	Food	Amount Mean	of iron in Standard error	100 grams Number of samples	Confi- dence code ¹	AH-8 Item No. (1963)
		Milligrams				
	Grapefruit:					
82	Raw, pink or white	0.07	0.01	53	a**	1053
83	Juice pack	.21	.05	5	c*	
84	Sirup pack, light	.40	.03	14	b*	1070
85	Unsweetened	.20	.02	17	p**	1071
86	SweetenedGrapes:	.36	.03	16	b*	1072
87	Raw, European	.26	.02	44	a**	1085
88	Canned, sirup pack, heavy	.94	(²)	2	b**	1087
89	Juice, canned, purple Mandarin oranges:	.24	.03	17	a**	1088
90	RawCanned:	.10	.01	16	C*	2262
91	Juice pack	.26	(²)	2	c*	
92	Sirup pack, light Mixed fruit:	.37	.03	32	b*	
93	Canned, sirup pack, heavy	.36	.03	16	b*	
94	Frozen, sweetened	.28	.04	3	c*	
95	Muskmelon, cantaloup, raw	.21	.02	77	a**	1358
96	Nectarines, raw	.15	.01	38	Ъ	1374
97	Raw	.08	.004	91	b**	1420
98	Juice, frozen, reconstituted	.11	.003	457	c*	1437
99	Papayas, raw	.10	.02	20	a**	1471
100	Raw	.11	.01	84	a**	1479
101	Water pack	.32	.04	16	Ъ	1480
102	Juice packSirup pack:	.27	.03	17	Ъ	1481
103	Light	.36	.03	19	b*	1482
104	Heavy	.27	.01	316	а	1483
105	Dried, sulfured	4.1	.12	24	Ъ	1487
106	Frozen, sweetened	.37	.07	3	c*	1490
107	Nectar, canned Pears:	.19	.04	5	b**	1491
108	RawCanned:	.25	.01	42	a**	1502
109	Water pack	.21	.02	12	C*	1504
110	Juice pack	.29	.06	7	b**	1505
111	Light	.28	.01	20	b*	1506
112	Heavy	.22	.01	105	a	1507

See footnotes at end of table.

TABLE 3.--IRON CONTENT OF EDIBLE PORTION OF FOOD--Continued

Item No.	Food	Amount Mean	of iron in Standard error	Number of samples	Confi- dence code ¹	AH-8 Item No. (1963)
		Milligrams				
113	Dried, sulfured	2.1	0.09	18	Ъ	1509
114	Nectar, canned Pineapples:	.26	.03	8	a	1512
115	RawCanned:	.37	.02	15	а	1611
116	Juice pack	.28	.01	676	a**	1614
117	Sirup pack, heavy	.38	.01	185	a**	1616
118	Juice, canned, unsweetened	.26	.01	77	a**	1619
119	Plums, canned, juice pack Prunes:	•34	.04	6	c*	
120	Canned, sirup pack, heavy	.41	.03	20	b**	
121	Dried	2.5	.07	193	a**	1818
122	Juice, canned	1.2	•20	30	a**	1821
123	Seeded (Muscat) Thompson seedless:	2.6	•29	34	b**	
124	Golden	1.8	.03	22	b*	
125	Natural	2.1	•09	51	a**	1846
126	Raw	.57	.03	19	Ъ	1849
127	Frozen, sweetened	.65	.10	3	c*	1852
128	Rhubarb, raw	.22	.02	33	b**	1865
129	RawFrozen:	.38	•04	84	a**	2217
130	Unsweetened	.75	.28	6	b**	
131 132	Sweetened, sliced Tropical fruit salad,	•59	.04	18	a**	2219
	Canned, sirup pack, heavy	•52	.02	25	b*	
133	Watermelon, raw	.17	.01	45	a**	2424
	GRAIN PRODUCTS					
134	Barley, pearled, uncooked	2.1	.15	5	C*	145, 146
135	Bulgur (parboiled wheat), uncooked	5.6	.74	14	a**	497, 498, 499
	Cornmeal:			0.4.0	4 .4.	000
136 137	Whole ground	1.8	.05 .07	243 144	b* b	883 885 1298 1377

Continued--

	Food	Mean	Standard error	Number of samples	Confi- dence code ¹	AH-8 Item No. (1963)
		Milligrams				
138	Pasta, enriched, uncooked	3.9	0.25	18	a	1298, 1377, 2157
139	Popcorn, popped with oil Rice, uncooked:	3.0	(²)	2	Ъ*	1655
140	Brown	1.8	.23	3	а	1869
141 142	Unenriched	1.2 ·4.6	•20 •27	6 18	b a**	1877 1871, 1873, 1875
	Spaghetti. See Pasta, item 138. Wheat:					2157
143	Bran, crude	10.8	.66	16	а	2446
144 145 146	Whole	4.3 3.5 ⁴ 5.6	.10 .06 .48	21 79 13	a a a**	2435 2439 2447
	LAMB, VEAL, AND GAME					
147	Frog, leg, raw	1.9	(2)	2	C *	1020
148 149 150	Chop, lean meat, broiled Leg, lean meat, roasted Veal, lean meat, raw	1.8 2.2 .67	.11 .12 .11	27 29 4	b* b* b**	1217 1187
	LEGUME PRODUCTS 3					
	Beans, common: Boiled:					
151 152 153 154 155 156	Great northern	2.3 3.6 2.6 2.4 3.0 3.0	.17 .27 .08 .17 .16	16 14 12 5 16 5	a** a b* a a	 155
157 158 159	Canned: Kidney Pinto Red.	1.3 2.0 1.5	.04 .83 .05	38 4 11	b a** c*	 161
	White, with sauce:	1.5	.03	52	a	

TABLE 3.--IRON CONTENT OF EDIBLE PORTION OF FOOD--Continued

Item No.		Amount Mean	of iron in Standard error	Number of samples	Confi- dence code ¹	AH-8 Item No. (1963)
-		Milligrams	are dallar villa vil			
162	With frankfurters	1.7	0.09	26	Ъ	
163	With pork Beans, lima: Boiled:	1.6	•02	182	a	
164	Baby	2.6	.06	13	а	
165	Large	3.1	.16	14	a**	
166	Canned, large	2.2	.16	6	c**	
167	Broad beans, canned	1.0	(²)	1	b*	
168	Boiled	3.0	.17	17	a**	
169	Canned	1.2	(²)	2	b	
170	Chili, with beans, canned Cowpeas (blackeye peas):	1.9	.09	16	c*	756
171	Boiled	3.8	.36	17	a**	904
172	Plain	.78	.16	3	a**	
173	With pork	1.4	•05	3	c*	
174	Lentils, boiled, Peanuts:	3.6	•24	16	a**	1254
175	Boiled	1.0	(²)	1	b*	1494
176	Roasted with skins	2.1	.17	9	a	1495
177	Peanut butter, chunky or smooth	1.7	.02	164	a**	
178	Peas, split, boiled	1.1	.03	12	Ъ*	1533
179	Soybeans, fermented product, miso	3.6	(²)	2	b**	2142
180	Soybean curd (tofu)	1.8	.25	3	Ъ	2145
	MIXED DISHES ³					
181	Beef stew, canned	1.3	.21	8	a**	372
182	Beef	1.4	.08	17	a**	
183	Fish	.73	.04	12	a	
184	Poultry	1.1	.06	16	a	
185	Shrimp Fast foods:	1.4	(2)	2	b	
186	Cheeseburger	2.2	•22	5	a**	
187	Fish sandwich	1.0	.15	4	a**	
188	Hamburger	2.6	.27	5	a	
189	Hot dog	1.6	(²)	2	b	
190	Beef	1.5	.47	3	b**	383
191	Poultry	.84	.14	6	b**	
TZT			(²)			

Continued--

Item No.	Food	Amount Mean	of iron in Standard error	100 grams Number of samples	Confi- dence code ¹	AH-8 Item No. (1963)
		Milligrams				
	NUTS AND SEEDS					
193	Almonds, dried	4.5	0.09	62	a	8
194	Cashew nuts	6.4	.21	14	b**	628
195	Coconut meat, dried, unsweetened.	5.0	()	2	b**	790
196	Filberts (hazelnuts)	8.1	()	2	Ъ	1008
197	Pecans	2.6	.37	14	a**	1536
198	Pistachio nuts	6.7	•55	13	a**	1626
199	Pumpkin seed kernels, dry	10.0	.77	4	С	1833
200	Sunflower seed kernels, dry	4.5	1.4	3	C**	2236
201	Walnuts, English	3.5	•54	8	a**	2421
	PORK PRODUCTS					
	Cured:					
202 203	Bacon, fried	1.7	.14	34	a**	126
	roastedFresh:	1.3	•05	3	b*	
204	Leg, shoulder, or sirloin,					
	lean, roasted	1.2	.03	59	Ъ	
205	Liver, raw	22.1	1.6	33	a**	1273
206	Loin chop, lean, broiled	.81	.04	12	Ъ	1720
	SUGARS AND SWEETS					
	Candy, chocolate:					
207	Milk, plain	1.2	.08	7	a	587
208	Semi-sweet	2.8	.29	6	a**	585
209	Sweet	2.2	.20	3	b*	586
210	Chocolate, baking	5.5	.62	5	a	759
211	Chocolate sirup, thin type	2.6	•20	15	a**	760
	Cocoa, dry powder:					700
212	Dutch	15.0	2.6	6	b*	782, 784,
						786
213	Plain	11.1	1.4	6	b*	781, 783,
	Molasses, cane:					785
214	Blackstrap	25.2	3.2	4	Ъ	1341
215	Light	5.8	1.1	4	b**	1339
					Cont	inued

See footnotes at end of table.

Item No.	Food	Amount Mean	of iron in Standard error	100 grams Number of samples	Confi- dence code ¹	AH-8 Item No. (1963)
				- July 100		(1703)
		Milligrams				
	Sirups:					
216	Cane	3.4	(²)	2	b	2048
217	Dark	1.0	(²)	1	c*	
218	Light	.1	(2)	1	C*	
219	Maple	1.2	(2)	1	c*	2049
220	Sorghum	14.0	5.1	4	C*	2050
221 222	Corn sirup with sugar Cane and corn sirups	.10	(2)	1	C*	
lus lus lus	with sugar	2.5	(²)	1	c*	
223	Cane, corn, and maple sirups.	2.7	(²)	1	c*	
224	Sugar, beet or cane, brown	2.7	.29	5	a**	2229
	VEGETABLE PRODUCTS ³					
225	Asparagus, green, canned	•54	.03	171	a**	48
0.0.6	Beans, canned:	1 (0.5	107	a**	1.00
226	Lima	1.6	.05	137		166
227	Snap, green or yellow	.88	•04	1426	a**	185,
220	Donn and (man)	1 6	•32	4	a**	197 180
228	Bean sprouts (mung), raw Beets:	1.6		4		
229	Raw	.91		5	a**	384
230	Canned	.67	.03	190	a**	386
231	Broccoli, raw	1.1	.30	9	а	483
232	Brussels sprouts, raw	1.4	(²)	2	b**	489
233	Cabbage, common varieties, raw Carrots:	•56	.15	29	a**	512
234	Raw	.51	.02	217	a**	619
235	Canned	.63	.03	157	a**	621
236	Cauliflower, raw	.58		4	a**	630
237	Celery, Pascal type, raw	.48		19	a**	637
238	Collards, frozen	1.1	.11	13	a**	810
239	Canned, regular pack	.33	.01	229	a**	847, 848, 849
240	Frozen, kernels, cut off cob	•45	•05	16	a**	856
241	Cucumber, pared, raw	.30		8		943
242	Eggplant, raw	•54		4	a b**	986
243	Kale, leaves only, raw	1.7	(²)	1	c*	1153
244	Lettuce, raw: Iceberg	•57	.19	72	a**	1258
245	Romaine	1.1	(²)	1	a** c*	1257
See	footnotes at end of table.				Cont	inued

Item No.	Food	Amount Mean	of iron in Standard error	100 grams Number of samples	Confi- dence code ¹	AH-8 Item No. (1963)
		Milligrams				
	Mushrooms:					
246	Raw	1.7	.20	26	a**	1354
247	Canned, drained solids	.79	.08	6	b**	
248	Mustard greens, raw	1.5	(²)	2	b**	1366
249	Okra, raw	.80	(²)	1	c*	1402
250	Mature, raw	.36	.14	44	a**	1412
251	Young green, bulb and					
	entire top, raw	1.9	1.1	3	a**	1415
252	Parsley, raw	1.6	(²)	2	Ъ	1472
253	Parsnips, raw	.58	(²)	2	Ъ	1473
254	Peas, sweet, canned	1.1	.02	478	a**	1523
255	Peas and carrots, canned	.74	•02	165	a	
256	Peas and onions, frozen	1.5	(²)	2	c*	
257	Peppers, sweet, raw	1.3	.82	5	a**	1545
	Potatoes, pared:					
258	Raw	.76	.04	84	a**	1785
259	Boiled	•35	.08	6	c*	1788
260	Canned	1.0	.13	30	Ъ	1796
261	Pumpkin, canned	1.4	.10	106	a**	1832
262	Sauerkraut, canned Spinach:	1.5	.09	173	Ъ	1977
263	Raw	2.7	•52	10	a**	2169
264	Canned	1.7	•04	230	a	2171
265	Summer varieties	.43	.03	11	b	2191
266	Winter varieties Sweetpotatoes:	•54	.07	5	C*	2199
267	All commercial varieties, raw	.59	.03	40	a**	2246
268	Canned, sirup pack, light Tomatoes:	.89	•05	70	b*	2252
269	Raw	.48	.01	141	a**	2282
270	Stewed	.61	.02	229	a**	
271	Whole	.60	•02	405	a**	2284
272	Tomato juice, canned	•55	.02	175	a**	2288
273	Tomato paste	3.0	.23	330	a**	2295
274	Tomato sauce, plain Turnip greens:	.72	.01	525	a**	
275	Raw	1.1	(²)	1	c*	2354
276	Canned	1.5	.13	5	c*	2357
2 / U	Vegetable juice, canned	.44	.03	13		2396

¹See p. 3 for explanation of confidence code.

²No standard error is given when the number of samples is two or less.

³Data for all canned items are for solids and liquid unless otherwise specified.

⁴This value is derived from analytical data. The new enrichment standard for flour (effective July 1983) will be 4.4 mg iron per 100 g.

LITERATURE CITED

- (1) Pao, E. M., and Mickle, S.J.

 1981 Problem nutrients in the United States. Food
 Technology. 35:58-69.
- (2) National Academy of Sciences National Research Council.
 1980 Recommended dietary allowances. 9th ed., rev.,
 185 pp. Washington, D.C.
- (3) Watt, B. K., and Merrill, A. L. 1963 Composition of foods... raw, processed, prepared. U.S. Dept. Agr. Agr. Handb. 8 (rev.), 190 pp.
- (4) Consumer and Food Economics Institute.
 1976 Composition of foods: Dairy and egg products; raw,
 processed, prepared. U.S. Dept. Agr. Agr. Handb.
 8-1, 157 pp.
- 1977 Composition of foods: Spices and herbs; raw, processed, prepared. U.S. Dept. Agr. Agr. Handb. 8-2, 51 pp.
- 1978 Composition of foods: Baby foods; raw, processed prepared. U.S. Dept. Agr. Agr. Handb. 8-3, 231 pp.
- 1979 Composition of foods: Fats and oils; raw, processed, prepared. U.S. Dept. Agr. Agr. Handb. 8-4, 142 pp.
- 1979 Composition of foods: Poultry products; raw, processed, prepared. U.S. Dept. Agr. Agr. Handb. 8-5, 330 pp.
- 1980 Composition of foods: Soups, sauces, and gravies; raw, processed, prepared. U.S. Dept. Agr. Agr. Handb. 8-6, 228 pp.
- (10) Consumer Nutrition Center.

 1980 Composition of foods: Sausages and luncheon meats;
 raw, processed, prepared. U.S. Dept. Agr. Agr. Handb.
 8-7, 92 pp.
- 1982 Composition of foods: Breakfast Cereals: raw, processed, prepared. U.S. Dept. Agr. Agr. Handb. 8-8, 162 pp.
- (12) Code of Federal Regulations.
 1980 Title 21, Section 101.9.
- (13) Monsen, E. R., Hallberg, L., Layrisse, M., Hegsted, D. M., Cook, J. D., Mertz, W., and Finch, C. A. 1978 Estimation of available dietary iron. Am. J. Clin. Nutr. 31:134-141.

United States Department of Agriculture Human Nutrition Information Service Washington, D.C. 20250

OFFICIAL BUSINESS
Penalty for Private Use, \$300

Postage and Fees Paid U.S. Department of Agriculture AGR -101

